

ORBIFOLD SPLICE QUOTIENTS AND LOG COVERS OF SURFACE PAIRS

WALTER D. NEUMANN AND JONATHAN WAHL

ABSTRACT. A three-dimensional orbifold (Σ, γ_i, n_i) , where Σ is a rational homology sphere, has a universal abelian orbifold covering, whose covering group is the first orbifold homology. A singular pair (X, C) , where X is a normal surface singularity with \mathbb{Q} HS link and C is a Weil divisor, gives rise on its boundary to an orbifold. One studies the preceding orbifold notions in the algebro-geometric setting, proving the existence of the universal abelian log cover of a pair. A key theorem computes the orbifold homology from an appropriate resolution of the pair. In analogy with the case where C is empty and one considers the universal abelian cover, under certain conditions on a resolution graph one can construct pairs and their universal abelian log covers. Such pairs are called orbifold splice quotients.

Let $(X, 0)$ be the germ of a normal complex surface singularity whose link Σ is a rational homology sphere (\mathbb{Q} HS). The topology of Σ is determined from any good resolution

$$(\tilde{X}, E) \rightarrow (X, 0)$$

by its weighted dual resolution graph Γ , which is a tree. The discriminant group $D(\Gamma)$, the cokernel of the intersection matrix $(E_i \cdot E_j)$, is isomorphic to the first homology group of Σ . The universal abelian cover (UAC) $\Sigma' \rightarrow \Sigma$ extends to a “cover” $(X', 0) \rightarrow (X, 0)$ of singularities, also called the UAC. These covers are quotients by an action of $D(\Gamma)$.

It was shown in [7] that under some mild conditions on a graph Γ with t ends (“semigroup and congruence conditions”), one can construct

- (1) explicit classes of complete intersection singularities $(X', 0) \subset (\mathbb{C}^t, 0)$
- (2) a diagonal representation $D(\Gamma) \subset (\mathbb{C}^*)^t$ acting freely on each $X' - \{0\}$
- (3) $(X', 0) \rightarrow (X', 0)/D(\Gamma) \equiv (X, 0)$, the UAC of a singularity with graph Γ .

Such X , called “splice quotient singularities,” are thus explicit examples having a resolution with the given graph Γ . (See (6.1) below for a precise statement). For their role in singularity theory, see e.g. [4] and [5].

The “end-curve theorem” of [8] characterizes those singularities $(X, 0)$ which are splice quotients. The t ends of the graph Γ of the minimal good resolution correspond to t isotopy classes of knots in the link Σ . Splice quotient singularities are exactly those for which each such class is represented (up to a multiple) by the zero-set of a function on $(X, 0)$. In this way, one recovers the result ([10]) that rational singularities, and those minimally elliptic singularities with \mathbb{Q} HS link, are splice quotients.

Rather than consider only covering spaces of a three-manifold Σ , it is natural to consider *orbifold covers*. Given a class of r knots $\{\gamma_i\}$ in Σ , with multiplicities $n_i \geq 1$, consider “covers” which are allowed to branch over these knots, with ramification indices bounded by the n_i . (Compare with looking at extensions of a number field allowed to ramify over some finite set of primes.) In Section 1 this notion is discussed and related to usual orbifold language, including

2000 *Mathematics Subject Classification.* 32S50, 14J17, 57M25, 57N10.

Key words and phrases. surface singularity, splice quotient singularity, orbifold homology, rational homology sphere, singular pair, abelian cover.

Neumann’s research supported under NSF grant no. DMS-1608600.

the *universal abelian orbifold cover* (UAOC) $\Sigma'' \rightarrow \Sigma$ and its covering group, the *(first) orbifold homology group* $H_1^{orb}(\Sigma)$. When Σ is a QHS, Proposition 1.3 indicates how to compute $H_1^{orb}(\Sigma)$, and proves it is an extension of the usual first homology group of Σ by a product of cyclic groups of order n_i . Abelian orbifold covers are classified by quotients of $H_1^{orb}(\Sigma)$.


There is an analogous notion of cover for a singular pair (X, C) . Here $(X, 0)$ is again a normal surface singularity with link Σ , and $C = \sum_{i=1}^r n_i C_i$ is a sum of irreducible Weil divisors $\{C_i\}$. By intersecting with a neighborhood boundary, a pair (X, C) gives rise to Σ and the orbifold data of knots γ_i and weights n_i . A *log cover* of the pair is a finite covering of X branched only over the C_i , with ramification bounded by n_i . Basic results about branched cyclic covers are given in Section 2.

For (X, C) with QHS link, Theorem 3.1 proves the existence of the *universal abelian log cover*, or *UALC*, a log cover of pairs $(X'', C'') \rightarrow (X, C)$ which corresponds to the UAOC on the boundary. Note that the C_i need not be \mathbb{Q} -Cartier.

From the graph of any good resolution of $(X, 0)$, one can compute the covering group $D(\Gamma)$ of the UAC $(X', 0) \rightarrow (X, 0)$. To find the covering group of the UALC of a pair (X, C) , consider the smallest good resolution $(\tilde{X}, E) \rightarrow (X, 0)$ for which the proper transform \tilde{C}_i of each C_i intersects transversally a leaf (or end) E_i of E , and each leaf intersects at most one \tilde{C}_i . Such leaves are called *special*, and $\tilde{X} \equiv \tilde{X}_C$ is the *minimal orbifold resolution* of (X, C) . Denote the corresponding graph Γ_C ; it is quasi-minimal (see (6.1) below). Now let Γ_C^* be the same graph, but decorated by the placement of an arrow and a weight n_i at each special leaf E_i . Finally, let $D(\Gamma_C^*)$ be the cokernel of the matrix $(E_i \cdot E_j)$ modified by multiplying each row corresponding to a special leaf E_i by n_i . One can easily prove there is a short exact sequence

$$0 \rightarrow \oplus \mathbb{Z}/(n_i) \rightarrow D(\Gamma_C^*) \rightarrow D(\Gamma_C) \rightarrow 0.$$

Example. If $X = \mathbb{C}^2$ and $C = \{x^p y^q (x - y)^r (y^2 - x^3)^s = 0\}$, the diagram for Γ_C^* is



One finds Γ_C by removing the arrows and p, q, r, s . $D(\Gamma_C^*)$ is the direct sum of 4 cyclic groups, of orders p, q, r, s respectively.

The first important result is proved in (4.3):

Theorem 1. $D(\Gamma_C^*)$, the cokernel of the modified intersection matrix above, is isomorphic to $H_1^{orb}(\Sigma)$, and is the covering group of the UALC of (X, C) .

According to [7], a graph Γ with t ends gives a faithful diagonal representation $D(\Gamma) \hookrightarrow (\mathbb{C}^*)^t$. In Proposition 5.3, we show there is a representation $D(\Gamma^*) \hookrightarrow (\mathbb{C}^*)^t$ compatible with that of $D(\Gamma)$ via a map $N : (\mathbb{C}^*)^t \rightarrow (\mathbb{C}^*)^t$ which raises coordinates to the power given by the weight at the corresponding special end of Γ^* , and leaves other entries fixed.

Our purpose is to define what it means for a pair (X, C) to be an *orbifold splice quotient*. The graphs Γ_C and Γ_C^* depends only on the boundary topology of the corresponding orbifold. In (6.5) we have

Theorem 2. *Let Γ be a quasi-minimal graph with t ends and Γ^* a decorated version, assigning a weight to the special ends. Let $D(\Gamma^*) \hookrightarrow (\mathbb{C}^*)^t$ be the corresponding representation. If Γ satisfies the semigroup and congruence conditions, then*

- (1) *there are isolated complete intersection singularities $(X'', 0) \subset (\mathbb{C}^t, 0)$*
- (2) *on each $(X'', 0)$, there is a Cartier divisor C'' so that $D(\Gamma^*)$ acts on the pair (X'', C'') , freely off the support of C''*
- (3) *each quotient $(X'', C'') \rightarrow (X'', C'')/D(\Gamma^*) \cong (X, C)$ is the UALC*
- (4) *the orbifold data associated to (X, C) is Γ, Γ^* .*

Definition. A pair (X, C) in the above Theorem is called an *orbifold splice quotient*.

We outline the construction and proof. Assign a variable x_i to every end of Γ . By the semigroup and congruence hypotheses and the major Theorem 7.2 of [7], one may consider a set of $D(\Gamma)$ -invariant splice-equations $\{f_j(x_i) = 0\}$, giving an isolated complete intersection singularity $(X', 0) \subset (\mathbb{C}^t, 0)$, which is the UAC of the quotient $X \equiv X'/D(\Gamma)$. The Cartier divisor $\{x_i = 0\}$ on X' is a reduced curve; some power of x_i is $D(\Gamma)$ -invariant, and $x_i^{\det \Gamma}$ provides a Cartier divisor on X whose reduction is an irreducible \mathbb{Q} -Cartier divisor. In particular, for each special end of Γ and corresponding x_i , define the \mathbb{Q} -Cartier divisor C_i as the reduced image of $\{x_i = 0\}$ on $(X, 0)$.

Next, replace a variable x_i corresponding to a special leaf by $z_i^{n_i}$. Consider the new complete intersection singularity $(X'', 0) \subset (\mathbb{C}^t, 0)$ defined by $\{f_j(z_i^{n_i}) = 0\}$. This is the inverse image of $(X', 0)$ under the aforementioned power map $N : \mathbb{C}^t \rightarrow \mathbb{C}^t$ which raises to the n_i^{th} power in the i^{th} special entry, and is the identity on others. We prove that $(X'', 0)$ is an isolated complete intersection singularity. We also show (5.3) $D(\Gamma^*)$ is the inverse image under N of $D(\Gamma)$, hence acts on $(X'', 0)$. Letting C'' be the sum of the divisors $z_i = 0$ for the special leaves, the result will follow.

There is an analogue of the End-Curve Theorem of [8] for orbifold splice quotients. For a given pair (X, C) , consider the minimal orbifold resolution $(\tilde{X}_C, E) \rightarrow (X, 0)$ and the corresponding graph Γ_C , with the special ends noted. In (6.8) we prove:

Theorem 3. *(Orbifold End-Curve Theorem) A pair $(X, \sum n_i C_i)$ is an orbifold splice quotient if and only each C_i is \mathbb{Q} -Cartier, and for every non-special end of the graph Γ_C , there is a function on $(X, 0)$ whose zero-set on \tilde{X}_C cuts out a smooth curve transversal to the end.*

It follows from the definition that an orbifold splice quotient is a splice quotient for which each C_i is \mathbb{Q} -Cartier. We do not know whether the converse is true; we suspect not. However, in (6.9) and (6.10) we have the following:

Corollary. *If $(X, 0)$ has a rational singularity, then every pair (X, C) is an orbifold splice quotient.*

Corollary. *Let (X, C) be a pair for which every non-special end of Γ_C comes from an end of the minimal good resolution. Then (X, C) is an orbifold splice quotient if and only if $(X, 0)$ is a splice quotient and every C_i is \mathbb{Q} -Cartier.*

The paper is organized as follows: In Section 1 we recall the basics of orbifolds in dimension three, the universal orbifold covering and fundamental group, and orbifold homology. Section 2 introduces singular pairs and their log covers, with focus on cyclic covers. From now on, one assumes the link is a \mathbb{Q} HS. Section 3 gives an algebro-geometric construction of the universal abelian log cover of a pair. The orbifold homology is computed in Section 4 from a plumbing diagram; the crucial representation of the orbifold discriminant group occurs in Section 5. The

definition and construction of orbifold splice quotients appears in Section 6, with an example presented in Section 7.

We are grateful to Eduard Looijenga and Helge Møller Pedersen for assistance with proving Proposition 4.4.

1. ORBIFOLDS

For the special purposes of this paper, an *orbifold* will be a compact oriented 3-manifold Σ , plus a disjoint set of embedded circles (or “knots”) $\gamma_1, \dots, \gamma_r \subset \Sigma$, to which are associated positive integers n_1, \dots, n_r (the *orbifold weights*). While this is not the standard definition of orbifold (e.g., [5]), it will be useful in considering branched covers with restricted ramification. See Example 1.2 below.

Definition 1.1. A (finite) *orbifold covering* consists of another orbifold $(\Sigma', \gamma'_j, n'_j)$ and a finite map $f : \Sigma' \rightarrow \Sigma$, with the following properties:

- (1) the inverse image of a knot γ_i consists of various γ'_{i_j} , with n'_{i_j} dividing n_i
- (2) $f^{-1}(\cup \gamma_i) = \cup \gamma'_{i_j}$, and f is a covering map off this set
- (3) locally around γ'_{i_j} , f maps to a neighborhood of γ_i as an orbifold quotient of order n_i/n'_{i_j} .

The last statement means that if the pairs (Σ, γ_i) and (Σ', γ'_{i_j}) are given locally as $(\mathbb{R}^3, z\text{-axis})$, then the map f is given by dividing out by the group of rotations of order n_i/n'_{i_j} around the z -axis. Since f is a covering map of some degree d off the γ'_{i_j} , it follows that for each i ,

$$\sum_j n_i/n'_{i_j} = d.$$

More precisely, the orbifolds we are considering could be described as “locally cyclic orientable 3-orbifolds” since their local structure is always $D^3/(\mathbb{Z}/(n))$ for some n (D^3 a three-disk). We restrict to cyclic groups since all our 3-manifolds are boundaries of 4-manifolds with complex structure.

Example 1.2. Let M' be a compact oriented 3-manifold on which a finite group G acts smoothly and faithfully, preserving orientation, and with the property that the isotropy group G_x at any point x is cyclic. We consider $M' \rightarrow M \equiv M'/G$ as an orbifold covering. The knots $\gamma_i \subset M$ are the images of the components of the set of points on M' with non-trivial isotropy, and the weight n_i is the order of this isotropy group. The γ'_{i_j} are the inverse images of the γ_i , and the $n'_{i_j} = 1$.

From now on we will often use the notation Σ both for the orbifold and its underlying space when what is meant is clear from context. Most of the definitions here and below are due to Thurston (e.g., Chapter 13 of his notes [14]).

An orbifold has an *orbifold fundamental group* $\pi_1^{orb}(\Sigma)$, obtained by quotienting $\pi_1(\Sigma - \bigcup_i \gamma_i)$ by the relations $\mu_i^{n_i} = 1$, where μ_i is represented by the boundary of a small transverse disk to γ_i . This can also be described as $\pi_1^{orb}(\Sigma) = \pi_1(\Sigma')$, where Σ' is the result of replacing a $D^2 \times S^1$ neighborhood of each γ_i by $K(\mathbb{Z}/(n_i), 1) \times S^1$, glued so that μ_i represents a generator of $\pi_1(K(\mathbb{Z}/(n_i), 1))$. (Here $K(\mathbb{Z}/(n_i), 1)$ is an Eilenberg-MacLane complex—a CW-complex with fundamental group $\mathbb{Z}/(n_i)$ and contractible universal cover.)

The Galois correspondence between covering spaces and subgroups of the fundamental group extends to orbifolds, using orbifold covers and orbifold fundamental group (see e.g., [6]). In

particular, any orbifold has a *universal abelian orbifold cover*, or *UAOC*, classified by the commutator subgroup of $\pi_1^{orb}(\Sigma)$. Its covering transformation group is the abelianization $\pi_1^{orb}(\Sigma)^{ab}$ of $\pi_1^{orb}(\Sigma)$; this is also $H_1^{orb}(\Sigma; \mathbb{Z}) := H_1(\Sigma'; \mathbb{Z})$, the degree 1 orbifold homology of Σ (see [6], [12], and [13] for more on orbifold homology).

Not every orbifold is the quotient of a manifold by a finite group, as in Example 2.1; an example is $(S^2 \times S^1, \{p\} \times S^1, n)$ for $n > 1$. However, if Σ is a QHS we prove:

Proposition 1.3. *Let (Σ, γ_i, n_i) be an orbifold for which the underlying space Σ is a QHS. Then the UAOC $\tilde{\Sigma}^{ab}$ of Σ is a finite cover of Σ , and its covering group $G = H_1^{orb}(\Sigma; \mathbb{Z})$ sits in an exact sequence*

$$0 \rightarrow \mathbb{Z}/(n_1) \oplus \cdots \oplus \mathbb{Z}/(n_r) \rightarrow G \rightarrow H_1(\Sigma; \mathbb{Z}) \rightarrow 0.$$

The Proposition follows from two Lemmas.

Lemma 1.4. *If Σ is a QHS, there is an exact sequence*

$$0 \rightarrow \bigoplus_{i=1}^r \mathbb{Z} \rightarrow H_1(\Sigma - \bigcup_i \gamma_i) \rightarrow H_1(\Sigma) \rightarrow 0,$$

where the first map is dotting with (μ_1, \dots, μ_r) .

Proof. The long-exact sequence of the pair $(\Sigma, \Sigma - \bigcup_i \gamma_i)$ gives

$$0 = H_2(\Sigma) \rightarrow H_2(\Sigma, \Sigma - \bigcup_i \gamma_i) \rightarrow H_1(\Sigma - \bigcup_i \gamma_i) \rightarrow H_1(\Sigma) \rightarrow H_1(\Sigma, \Sigma - \bigcup_i \gamma_i) = 0.$$

The first term is 0 since Σ is QHS. Let $T = \cup_i T_i$ be the union of small closed tubular neighborhoods of the γ_i , each one a closed 2-disk bundle over an S^1 ; let T° and T_i° denote their interiors. Then the inclusion $\Sigma - \bigcup_i \gamma_i \subset \Sigma - T^\circ$ induces an isomorphism $H_j(\Sigma, \Sigma - \bigcup_i \gamma_i) \cong H_j(\Sigma, \Sigma - T^\circ)$. By excision (removing $\Sigma - T$), one has

$$H_j(\Sigma, \Sigma - T^\circ) \cong H_j(T, T - T^\circ) = H_j(T, \partial T) = \bigoplus_i H_j(T_i, \partial T_i).$$

But

$$H_2(T_i, \partial T_i) = \ker H_1(\partial T_i) \rightarrow H_1(T_i)$$

is the free \mathbb{Z} -module generated by μ_i , while $H_1(T_i, \partial T_i) = 0$, whence the result. \square

Lemma 1.5. $H_1^{orb}(\Sigma) \cong H_1(\Sigma - \bigcup_i \gamma_i) / (n_1 \mu_1, \dots, n_r \mu_r)$.

Proof. If G is a group, N a normal subgroup, then abelianizing G/N is the same as abelianizing G and then modding out by the image of N in G_{ab} ; either method gives $G/G'N$, where $'$ denotes commutator subgroup. For, every element of the commutator subgroup is a product of commutators, so the map $G' \rightarrow (G/N)'$ is surjective. Apply now to $G = \pi_1(\Sigma - \bigcup_i \gamma_i)$ and N the normal subgroup generated by the relations $\mu_i^{n_i} = 1$. Then $G/N = \pi_1^{orb}(\Sigma)$, and abelianizing gives $H_1^{orb}(\Sigma)$; $G_{ab} = H_1(\Sigma - \bigcup_i \gamma_i)$, and the image of N is $(n_1 \mu_1, \dots, n_r \mu_r)$. \square

Remark 1.6. The Proposition shows one could construct the UAOC by first forming the UAC $\Sigma' \rightarrow \Sigma$, taking the inverse images of the γ_i 's, and then making a sequence $\tilde{\Sigma}^{ab} \rightarrow \Sigma'$ of cyclic branched covers over these knots. The orbifold weights on $\tilde{\Sigma}^{ab}$ are equal to 1.

2. COVERS OF SINGULAR PAIRS

The singularity version concerns *singular pairs* (X, C) . Here, $(X, 0)$ is the germ of a complex normal surface singularity (thus homeomorphic to the cone over its link Σ), and $C = \sum_{i=1}^r n_i C_i$ is a positive combination of reduced and irreducible Weil divisors. Intersecting with the boundary of a small ball neighborhood of the singular point in a smooth ambient space gives an orbifold link in a natural way.

The data of a singular pair (X, C) is more frequently encoded in the literature as a *log pair* (X, Δ_X) , with the *boundary of X* the effective \mathbb{Q} -divisor

$$\Delta_X = \sum_{i=1}^r (1/n_i) C_i.$$

Log pairs are essential notions in algebraic geometry (e.g. [2]), and $1/n_i$ could be replaced by any real number between 0 and 1. However, for this paper the singular pair $(X, \sum n_i C_i)$ is a more convenient way to display branching data for branched covers.

We denote the *support of C* by $|C| = \cup_{i=1}^r C_i$. A *map of pairs* $f : (X', C') \rightarrow (X, C)$ is a finite map of normal germs $f : X' \rightarrow X$ so that $f(|C'|) = |C|$.

Definition 2.1. A *log cover of pairs* $f : (X', C') \rightarrow (X, C)$ is a map of pairs with the following additional properties:

- (1) $f^{-1}(|C|) = |C'|$, and f is an unramified covering space off this set.
- (2) For $C = \sum_{i=1}^r n_i C_i$ and $f^* C_i = \sum_{j=1}^{s_i} m_{ij} C'_{ij}$ (as Weil divisors), one has $m_{ij} | n_i$, all j .
- (3) $C' = \sum_{i=1}^r \sum_{j=1}^{s_i} n_i / m_{ij} C'_{ij}$.

In other words, for a log cover of pairs, the map $f : X' \rightarrow X$ is ramified only over the C_i , and the ramification index m_{ij} on any curve above it divides n_i . (In particular, if $n_i = 1$, no ramification is allowed above C_i .)

Remark 2.2. Note that if a map of pairs satisfies (1), then the conditions (2) and (3) above are together equivalent to a relation on the boundary divisors, namely

$$f^*(\Delta_X) = \Delta_{X'}.$$

We note the easy

Lemma 2.3. *Let $f : X' \rightarrow X$ be a finite map of germs of normal surface singularities. If f is ramified off the singular points, then there are unique minimal divisors C' on X' and C on X so that $f : (X', C') \rightarrow (X, C)$ is a log cover of pairs.*

Proof. Let C_1, \dots, C_r be the irreducible components of the branch curve of f on X . Write $f^* C_i = \sum_{j=1}^{s_i} m_{ij} C'_{ij}$ as before. Define $n_i = \text{lcm}(m_{ij}, j = 1, \dots, s_i)$. Then set $C = \sum_{i=1}^r n_i C_i$ and $C' = \sum_{i=1}^r \sum_{j=1}^{s_i} n_i / m_{ij} C'_{ij}$. The result now follows. \square

A log cover of pairs $(X', C') \rightarrow (X, C)$ is said to be *abelian* (resp., *cyclic*) if an abelian (resp., cyclic) group G acts on X' with quotient X , permuting the curves lying above each C_i .

To make a cyclic cover, take the n th root of an appropriate function h on X , and normalize. Let A be the analytic local ring of X , $h \in m_A$ non-0. Then $T^n - h$ is an irreducible polynomial over the quotient field of A iff h is not a d th power, where $d|n$. One may verify this condition as follows: if the divisor (h) of zeroes of h is the sum of irreducible Weil divisors $\sum_{i=1}^r k_i C_i$, then for any $d > 1$ dividing n we have that $(1/d) \sum_{i=1}^r k_i C_i$ is *not* a principal divisor. In this case, adjoining an n th root of h and normalizing gives an n -cyclic cover of pairs $(X', C') \rightarrow (X, C)$, for

appropriate C and C' . Specifically, examining the cyclic behavior over each C_i and normalizing, we find that $m_{ij} = n/\gcd(n, k_i) = n_i$, so that

$$C = \sum_{i=1}^r (n/\gcd(n, k_i))C_i, \quad C' = \sum_{i=1}^r \sum_{j=1}^{\gcd(n, k_i)} C'_{ij}$$

is as in Lemma 3.2. If $n|k_i$, then the coefficient of C_i is 1, so that no branching occurs over this curve and it need not be part of C . We summarize in the

Lemma 2.4. *Let $(X, 0)$ be a normal germ, $h \in m_A$ a non-0 function with divisor of zeroes $\sum_{i=1}^r k_i C_i$. Suppose $n > 1$ is such that h is not a d th power, for any $d > 1$ which divides n . Then adjoining an n th root of h and normalizing gives an n -cyclic cover of pairs $(X', C') \rightarrow (X, C)$, with*

$$C = \sum' (n/(\gcd(n, k_i))C_i,$$

where $'$ means the sum is taken over those C_i for which n does not divide k_i .

We illustrate the last construction by asking whether there exists an n -cyclic cover totally ramified over a given irreducible Weil divisor. Recall that $\text{Cl}(X)$, the *divisor class group* of X , is the free abelian group generated by the irreducible Weil divisors on X (or equivalently height one prime ideals in $A = \mathcal{O}_{X,0}$), modulo principal divisors.

Proposition 2.5. *Let $(X, 0)$ be a normal surface singularity, C_1 an irreducible Weil divisor, $n > 1$ a positive integer.*

- (1) *There exists an n -cyclic cover of pairs $(X', C') \rightarrow (X, nC_1)$ with C' irreducible if and only if the class of C_1 in $\text{Cl}(X)$ is divisible by n .*
- (2) *Such a cover is unique exactly when the link of X is a QHS and the discriminant group has order prime to n .*

Proof. As explained above, suppose one has an n -cyclic cover obtained from a function whose zero divisor is $\sum_{i=1}^r k_i C_i$. It is branched exactly over those C_i for which n does not divide k_i . To have a cover as in (1) requires therefore that an $h \in A$ exist with $(h) = k_1 C_1 + nD$, where k_1 and n are relatively prime. If uk_1 is congruent to 1 mod n , we can use h^u with $(h^u) = C_1 + nD'$. Thus, the class of C_1 is divisible by n .

Conversely, if the class of C_1 is divisible by n , then for some h in the quotient field of A and divisor G , we can write

$$C_1 = nG + (h).$$

Writing $h = j/k$, with $j, k \in A$, we have that $(jk^{n-1}) = C_1 + nG'$, where $G' = (k) - G$. So, adjoining an n^{th} root of the regular function (jk^{n-1}) yields a cyclic cover which (by Lemma 2.4) is branched only over C_1 .

As for uniqueness, recall that adjoining n th roots of two elements h and h' gives the same field extension if and only if h/h' is an n th power. Suppose there is an effective non-Cartier divisor C so that nC is Cartier, with $(\alpha) = nC$. Then the Cartier divisors of h and $h\alpha$ differ by a multiple of n ; but extracting n th roots gives different field extensions. So, the uniqueness question is exactly whether there are non-trivial elements of order dividing n in the divisor class group. If the link of X is not a \mathbb{Q} -homology sphere, then the divisor class group contains a vector space modulo a free \mathbb{Z} -module, hence contains elements of all finite order. If the link is a QHS, then as recalled in the next section the divisor class group is a direct sum of a complex vector space plus the (finite) discriminant group. The result follows. \square

We illustrate the last proposition with the following

Example 2.6. Consider the A_1 singularity $X = \{(x, y, z) \in \mathbb{C}^3 : xz - y^2 = 0\}$, C_1 the line $\{y = z = 0\}$, and C_2 the irreducible Cartier divisor $\{z - x^2 = 0\}$. The link of $(X, 0)$ is the lens space $L(2, 1) = \mathbb{R}P^3$ with fundamental group $\mathbb{Z}/(2)$, and the divisor class group is cyclic of order 2, generated by the class of C_1 . The principal divisor $(z) = 2C_1$, and z is not a power of any element. We consider n th root constructions of z and $z(z - x^2)$.

- (1) For $n = 2k + 1$, the n th root of z gives $(X', C'_1) \rightarrow (X, (2k + 1)C_1)$. X' is another A_1 singularity $\{xv - u^2 = 0\}$, C'_1 is the line $\{u = v = 0\}$, and the coordinates are related by $y = uv^k, z = v^{2k+1}$. The group action on X' is given by $(x, u, v) \mapsto (x, \zeta u, \zeta^2 v)$, where ζ is a primitive $(2k + 1)$ st root of 1.
- (2) For $n = 2k$, the n th root of z gives $(\mathbb{C}^2, \{v = 0\}) \rightarrow (X, kC_1)$. A generator of the covering group acts via $(u, v) \mapsto (-u, \zeta v)$, where ζ is a primitive $(2k)$ th root of 1. Thus, $x = u^2, y = -uv^k, z = v^{2k}$. The cover factors through the UAC, given by taking the square root of z (the case $k = 1$).
- (3) The square roots of $z - x^2$ and $z(z - x^2)$ give (Lemma 2.4) non-isomorphic double covers of $(X, 2C_2)$. In the first case, the cover is obtained by adjoining T with $T^2 = z - x^2$, giving a D_4 singularity $y^2 = x(T^2 + x^2)$. For the second cover, one adjoins $S^2 = z(z - x^2)$; normalize via $P = Sx/y$, since $P^2 = x(z - x^2)$, and note also that $SP = y(z - x^2)$. This second cover, of embedding dimension 5, is an $8/3$ cyclic quotient singularity. Example 3.4 below gives the full story.

3. UNIVERSAL ABELIAN LOG COVER OF A PAIR

If $(X, 0)$ has \mathbb{Q} HS link Σ , then for topological reasons the UAC on the boundary extends to a UAC of $X - \{0\}$, so taking integral close of X in this extension gives what we call the UAC $(X', 0) \rightarrow (X, 0)$. One can also construct $(X', 0)$ directly using a covering of a good resolution of X , as in [9], (3.1).

If (X, C) is a singular pair for which X has \mathbb{Q} HS link Σ , one might in principle try to extend a cover of the corresponding orbifold associated to Σ to a branched cover of $X - \{0\}$ and then of X . It seems more natural to make a direct algebro-geometric construction. We have the following definition/theorem.

Theorem 3.1. *Suppose (X, C) is a singular pair for which X has \mathbb{Q} HS link Σ . Then there exists a universal abelian log cover (or UALC) $(X'', C'') \rightarrow (X, C)$, a cover of pairs which induces the UAOC on the boundary.*

Proof. Let $(\tilde{X}, E) \rightarrow (X, 0)$ be a good resolution, and \mathbb{E} the lattice generated by the irreducible exceptional curves E_i . Following Mumford [3], one has a natural direct sum decomposition

$$Cl(X, 0) \cong H^1(\tilde{X}, \mathcal{O}_{\tilde{X}}) \oplus \mathbb{E}^*/\mathbb{E},$$

the direct sum of a complex vector space of dimension $p_g(X)$ plus the finite *discriminant group* isomorphic to $H_1(\Sigma; \mathbb{Z})$, of order $|\det(E_i \cdot E_j)|$ (see [9], (2.3) for a nice discussion.) The universal abelian cover $g : (X', 0) \rightarrow (X, 0)$ is unramified off the singular point, with covering group \mathbb{E}^*/\mathbb{E} . We indicate how to form the UALC via a sequence of cyclic covers of X' .

Write $C = \sum n_i C_i$ (we may as well assume all $n_i > 1$). Since a vector space is a divisible group, we may write the class of C_i in $Cl(X)$ uniquely as $[C_i] = n_i [D_i] + t_i$, where t_i is a torsion divisor class. But the class of $g^*(t_i)$ is trivial, i.e., $g^*[C_i] - n_i g^*[D_i]$ is trivial in $Cl(X')$. Thus $g^*C_i - n_i g^*D_i$ is a principal divisor, given by a unique (up to scalar) function h'_i in the quotient field of $A := \mathcal{O}_{X', 0}$. Writing $h'_i = j_i/k_i$, with $j_i, k_i \in A$, we see that $h_i = j_i k_i^{n_i - 1} \in A$ is a regular function, defining an effective divisor of the form $g^*(C_i) + n_i F_i$, for some F_i . Since g is a covering off the singular point, $g^*(C_i)$ is a reduced divisor. So, one can form the cyclic

cover by adjoining an n_i -th root of h_i and normalizing; this depends only on h'_i . But the other $g^*(C_j)$ share no common components with each other, so remain reduced when pulled back to the cyclic cover given by a root of h_i . Thus we can repeat the cyclic quotient construction, and eventually obtain a series of cyclic covers branched to order n_j over every $g^*(C_j)$, and hence over C_j . The final local ring is the normalization obtained after adjoining to A the n_i^{th} root of h_i , all $i = 1, \dots, r$. \square

We summarize in the

Corollary 3.2. *Let $(X, C = \sum_{i=1}^r n_i C_i)$ be a singular pair for which X has \mathbb{Q} HS link Σ . Then the universal abelian log cover $(X'', C'') \rightarrow (X, C)$ is constructed by forming the universal abelian cover of X , and then taking cyclic covers of orders n_1 through n_r . The covering group G sits in a short exact sequence*

$$0 \rightarrow \mathbb{Z}/(n_1) \oplus \dots \oplus \mathbb{Z}/(n_r) \rightarrow G \rightarrow H_1(\Sigma; \mathbb{Z}) \rightarrow 0.$$

Example 3.3. In Example (2.6(2)) above, taking a fourth root of the function z gives the UALC of $(X, 2C_1)$; thus, the covering group is cyclic, so the exact sequence above is not split.

Example 3.4. In Example (2.(6.3)) above, the UALC of $(X, 2C_2)$ is obtained by first taking the UAC of X ; set $x = u^2, y = uv, z = v^2$, and the inverse image of C_2 is $v^2 - u^4 = 0$. Taking the double branched cover gives $\mathbb{C}[u, v, t]/t^2 - v^2 + u^4$ as UALC, an A_3 -singularity, with curve the two components of $t = v^2 - u^4 = 0$. The covering group is the Klein four-group $\mathbb{Z}/2 \times \mathbb{Z}/2$, generated by $(t, v, u) \mapsto (-t, v, u)$ and $(t, v, u) \mapsto (t, -v, -u)$. The invariants of the first involution is the UAC, while that of the second is the D_4 -singularity described as the “first case” in Example (2.(6.3)). The invariants of the other involution, which is $-I$ on the coordinates, gives the $8/3$ cyclic quotient; in the previous notation, $S = tv$ and $P = tu$.

Example 3.5. Consider the pair consisting of $(\mathbb{C}^2, 0)$ and $r \geq 3$ lines L_1, \dots, L_r through the origin, with multiplicities $n_1, \dots, n_r \geq 2$. The UALC of $(\mathbb{C}^2, \sum_{i=1}^r n_i L_i)$ is a Brieskorn complete intersection of type (n_1, \dots, n_r) . Specifically, for any $r-2$ distinct numbers a_1, \dots, a_{r-2} , consider the singularity $(Y, 0)$ defined by

$$x_i^{n_i} + x_{r-1}^{n_{r-1}} + a_i x_r^{n_r} = 0, \quad i = 1, \dots, r-2,$$

and the map $f : (Y, 0) \rightarrow (\mathbb{C}^2, 0)$ defined on the ring level by

$$x \mapsto x_r^{n_r}, \quad y \mapsto x_{r-1}^{n_{r-1}}.$$

Thus, $y + a_i x$ maps to $-x_i^{n_i}$, so the map is branched over the corresponding line with multiplicity $n_i, i = 1, \dots, r-2$. It is not hard to see that f is the UALC over these lines and the coordinate axes, and is the quotient by the group $\mathbb{Z}/(n_1) \oplus \dots \oplus \mathbb{Z}/(n_r)$ acting diagonally on the coordinates in \mathbb{C}^r .

4. ORBIFOLD HOMOLOGY GROUP FROM PLUMBING

As always, $(\tilde{X}, E) \rightarrow (X, 0)$ will denote a good resolution of a singularity with \mathbb{Q} HS link Σ , where $E = E_1 \cup \dots \cup E_m$. From the weighted resolution dual graph Γ , one forms the lattice $\mathbb{E} = \oplus \mathbb{Z} \cdot E_i$ and its dual \mathbb{E}^* , with dual basis e_1, \dots, e_m defined by $e_i(E_j) = \delta_{ij}$. The following is well-known:

Proposition 4.1. *$H_1(\Sigma)$ is isomorphic to the discriminant group $D(\Gamma) = \mathbb{E}^*/\mathbb{E}$, with generators e_1, \dots, e_m , modulo the relations*

$$\sum_{j=1}^m (E_i \cdot E_j) e_j = 0, \quad i = 1, 2, \dots, m.$$

This result follows from examining the exact sequence

$$H_2(\Sigma) = 0 \rightarrow H_2(\tilde{X}) \rightarrow H_2(\tilde{X}, \Sigma) \rightarrow H_1(\Sigma) \rightarrow H_1(\tilde{X}) = 0.$$

Note $H_2(\tilde{X}) = \oplus \mathbb{Z}[E_i] \cong \mathbb{E}$ and $H_2(\tilde{X}, \Sigma) \cong H^2(\tilde{X}) \cong H_2(\tilde{X})^* \cong \mathbb{E}^*$. More precisely, let $K_i \subset \Sigma$ be a meridian knot over E_i , i.e., the boundary of a complex disk A_i in \tilde{X} which is transversal to E_i . Then the classes $[A_i]$ form a basis of $H_2(\tilde{X}, \Sigma)$, dual to the $[E_i]$, and the image of $[A_i]$ in $H_1(\Sigma)$ is $[K_i]$.

The \mathbb{Q} -valued pairing on \mathbb{E}^* allows one to define $e_i \cdot e_j \in \mathbb{Q}$; by linear algebra, the matrix $(e_i \cdot e_j)$ is the inverse of the matrix $(E_i \cdot E_j)$. From ([8], (1.2) and (9.2)), one can calculate the topologically defined *linking number* ℓ of meridian knots on Σ :

Proposition 4.2. *The linking number satisfies*

$$\ell(K_i, K_j) = -e_i \cdot e_j.$$

A surface pair (X, C) gives rise to an orbifold (Σ, γ_i, n_i) on the boundary, and we will prove one can compute the orbifold homology $H_1^{orb}(\Sigma)$ from a *log resolution* of the pair. That is, we consider a good resolution $(\tilde{X}, E) \rightarrow (X, 0)$ for which the full inverse image of C has strong normal crossings. A proper transform \tilde{C}_i of C_i intersects one exceptional curve E_i transversally, so that its boundary γ_i is a meridian knot of type K_i over a point of E_i ; to it is associated an orbifold weight $n_i \geq 1$. For convenience, one can blow up further so that each exceptional curve intersects at most one \tilde{C}_i .

More generally, assume one is given a good resolution with exceptional curves E_1, \dots, E_m , and a meridian knot K_i and weight n_i associated to each E_i . This defines an orbifold structure (Σ, γ_i, n_i) on the link. The goal will be to prove the

Theorem 4.3. *For the orbifold (Σ, γ_i, n_i) , the orbifold homology $H_1^{orb}(\Sigma)$ is generated by e_1, \dots, e_m , modulo the relations*

$$\sum_{j=1}^m n_i (E_i \cdot E_j) e_j = 0, \quad i = 1, 2, \dots, m.$$

In other words, multiply the rows of the intersection matrix by the corresponding orbifold weight, and take the cokernel of the resulting non-symmetric matrix. Also, note that if $n_i = 1$, then K_i and γ_i are irrelevant.

By Lemma 1.5, to prove the Theorem one needs to find first the cohomology of $H_1(\Sigma - \bigcup \gamma_i)$. By Lemma 1.4 and the above discussion, one has as generators the classes of K_i and μ_j , and the complete set of relations must be of the form

$$(*) \quad \sum_{j=1}^m (E_i \cdot E_j) [K_j] = \sum_{j=1}^m n_{ij} [\mu_j], \quad i = 1, \dots, m,$$

for some integral matrix (n_{ij}) .

Note there is a natural orientation of the μ_j ; a small tubular neighborhood T_j of γ_j in Σ is a complex disk in E_j times γ_j , and μ_j is the boundary of a transverse disk.

Because of Lemma 1.5, Theorem 4.3 follows easily from the next Proposition. The proof is due to Helge Møller Pedersen, after which an alternate proof due to Eduard Looijenga will be outlined.

Proposition 4.4. *With the notation above, the relations in $H_1(\Sigma - \bigcup \gamma_i)$ are*

$$[\mu_i] + \sum_{j=1}^m (E_i \cdot E_j) [K_j] = 0, \quad i = 1, \dots, m.$$

Proof. We use linking numbers to relate the classes of the K_i and the μ_j .

Lemma 4.5. *Suppose K is a meridian as above, with $[K]$ of order d in $H_1(\Sigma)$. Then there is a closed oriented two-chain in $\Sigma - \bigcup \gamma_i$ whose boundary is homotopic to*

$$dK - \sum_{j=1}^m d\ell(K, \gamma_j)\mu_j.$$

Proof. There is a closed oriented 2-chain A in Σ whose boundary consists of d copies of K . We may modify A around the intersection points with all the γ_j so that there are $d|\ell(K, \gamma_j)|$ transverse intersection points of A with γ_j (itself a meridian curve). As $\ell(K, \gamma_j)$ is positive by Proposition 4.2, γ_j intersects A positively at these points. A small closed tubular neighborhood of γ_j intersects A in $d\ell(K, \gamma_j)$ small disks, and the boundary of each is homotopic to $+\mu_j$. Removing the interior of these disks from A therefore gives a closed 2-chain in $\Sigma - \bigcup \gamma_i$ with boundary $dK - \sum d\ell(K, \gamma_j)\mu_j$, as claimed. \square

Denoting by $\delta = |\det(E_i \cdot E_j)|$ the order of $H_1(\Sigma)$, we multiply by δ/d , and conclude the relations in $H_1(\Sigma - \bigcup \gamma_i)$

$$\delta[K_i] = \sum_{j=1}^m \delta\ell(K_i, K_j)[\mu_j], \quad i = 1, \dots, m.$$

Viewing as a matrix equation, left multiply both sides by $(E_i \cdot E_j)$, which by (4.2) is minus the inverse of $\ell(K_i, K_j)$; this gives

$$\delta \sum_{j=1}^m (E_i \cdot E_j)[K_j] = -\delta[\mu_i].$$

Substituting into (*) above yields

$$\delta \sum n_{ij}[\mu_j] = -\delta[\mu_i].$$

But the $[\mu_i]$ are \mathbb{Z} -independent in $H_1(\Sigma - \bigcup \gamma_i)$. Thus $n_{ij} = -\delta_{ij}$, so that

$$\sum_{j=1}^m (E_i \cdot E_j)[K_j] = -[\mu_i].$$

\square

Remark 4.6. Looijenga's approach is for each E_i to find the boundary of an appropriate closed oriented 2-chain in $\Sigma - \bigcup \gamma_i$. Let $b = -E_i \cdot E_i$.

Suppose γ_i lies over $r \in E_i$; p_1, \dots, p_t are intersection points of E_i with its neighbors; and q_1, \dots, q_b are any other distinct points of E_i . There is a meromorphic section s of the normal bundle $N \rightarrow E_i$, with simple poles at the q_k and no zeroes. Let E'_i result from removing from E_i the interiors of small disks centered at r , the p_j , and the q_k ; one now has the section $s : E'_i \rightarrow N$, which can be scaled into a section of an appropriate unit circle bundle $N' \subset N$. Now consider the images in N' of the boundaries of the removed disks.

As $N' \rightarrow E_i$ is locally $D \times S^1 \rightarrow D$, a fibre K_i is homotopic to the inverse image of the boundary of D (consider the family $(te^{i\theta}, e^{i\theta}), 0 \leq t \leq 1$). Since s has a simple pole at q_k , the image of the boundary of D has the opposite orientation as the inverse image (cf. $z \mapsto (z, 1/z) \mapsto (z, |z|/z)$). So the b circles contribute $-bK_i$ to the boundary of $s(E'_i)$. The circle around r contributes the knot μ_i to the boundary.

Since disks were removed from the intersections of E_i with other E_k , the 2-chain $s(E'_i)$ can be pushed up from N' minus these disks to $\Sigma - \bigcup \gamma_i$. Locally $E_i \cap E_k$ looks analytically like

$zw = 0$, and Σ can be viewed as $|zw| = \epsilon$. The boundary of the disk $\Delta_k = \{|z| \leq \delta, w = 0\}$ has boundary that can be lifted to $\{|z| = \delta, w = \epsilon/\delta\}$, which on Σ gives a meridian of E_k , of class K_k .

Therefore, the closed oriented 2-chain $s(E'_i) \subset \Sigma - \bigcup \gamma_i$ has boundary homotopic to

$$-bK_i + \mu_i + \sum_{k \neq i} (E_i \cdot E_k)K_k = \mu_i + \sum_k (E_i \cdot E_k)K_k.$$

This expression becomes zero in homology.

5. ACTION OF THE ORBIFOLD DISCRIMINANT GROUP

With (\tilde{X}, E) as before, suppose E_1, \dots, E_t are the ends (or *leaves*) of the graph Γ . Recall the diagonal representation of the discriminant group $D(\Gamma)$:

Proposition 5.1. ([7], (5.2), (5.3)). *There is a natural injection $D(\Gamma) \hookrightarrow (\mathbb{Q}/\mathbb{Z})^t$, given by $e \mapsto (e \cdot e_1, \dots, e \cdot e_t)$. Exponentiating $\mathbb{Q}/\mathbb{Z} \hookrightarrow \mathbb{C}^*$ via $r \mapsto \exp(2\pi ir)$, one has a faithful diagonal action of $D(\Gamma)$ on \mathbb{C}^t , where the entries are t -tuples of $\det(\Gamma)$ -th roots of unity.*

Given a pair (X, C) , there is a log resolution so that each \tilde{C}_i intersects an end E_i of the graph (cf. (6.4) below). For this orbifold situation, to those ends E_i we are given knots γ_i of type K_i plus a weight n_i ; assign a weight $n_j = 1$ to any other end. We represent the extra data by a decorated graph Γ^* , by adding to each end of Γ an arrow and the associated weight.

Definition 5.2. The orbifold discriminant group $D(\Gamma^*)$ is the group generated by e_1, \dots, e_m modulo the relations

$$\begin{aligned} \sum_{j=1}^m n_i(E_i \cdot E_j)e_j &= 0, \quad i = 1, 2, \dots, t. \\ \sum_{j=1}^m (E_i \cdot E_j)e_j &= 0, \quad i = t+1, \dots, m. \end{aligned}$$

According to Theorem 4.3, $D(\Gamma^*)$ is isomorphic to the orbifold homology group (all weights of interior E_j equal 1). Modding out the e_i 's by the further relations with all $n_i = 1$ (as in Proposition 4.1) gives the surjection $\Phi : D(\Gamma^*) \rightarrow D(\Gamma)$, with kernel the sum of cyclic groups of orders n_1 through n_t . This is the same as the general result in Proposition 1.3.

$D(\Gamma^*)$ has a natural diagonal representation compatible with that for $D(\Gamma)$, using the ‘‘power map’’ $N : (\mathbb{C}^*)^t \rightarrow (\mathbb{C}^*)^t$ given on points by

$$(a_1, \dots, a_t) \mapsto (a_1^{n_1}, \dots, a_t^{n_t}).$$

Proposition 5.3. *There is a natural injection $D(\Gamma^*) \hookrightarrow (\mathbb{C}^*)^t$ and a commutative diagram*

$$\begin{array}{ccc} D(\Gamma^*) & \hookrightarrow & (\mathbb{C}^*)^t \\ \downarrow \Phi & & \downarrow N \\ D(\Gamma) & \hookrightarrow & (\mathbb{C}^*)^t \end{array}$$

Proof. We claim there is a map $D(\Gamma^*) \rightarrow (\mathbb{Q}/\mathbb{Z})^t$ given by

$$e \mapsto ((e \cdot e_1)/n_1, (e \cdot e_2)/n_2, \dots, (e \cdot e_t)/n_t).$$

One needs to check that the relations defining $D(\Gamma^*)$ go to 0 in every entry of $(\mathbb{Q}/\mathbb{Z})^t$. But in the k th entry,


$$\sum_{j=1}^m n_i(E_i \cdot E_j)e_j \mapsto \sum_{j=1}^m n_i(E_i \cdot E_j)e_j \cdot e_k/n_k = n_i/n_k \sum_{j=1}^m (E_i \cdot E_j)e_j \cdot e_k = n_i/n_k \delta_{ik},$$

which equals 0 or 1, in either case 0 in \mathbb{Q}/\mathbb{Z} . After exponentiation, one gets a map to $(\mathbb{C}^*)^t$. This gives the commutative diagram asserted.

There remains to check injectivity in the top row, or equivalently injectivity of $\ker \Phi \rightarrow \ker N$. As these groups have the same order $n_1 \cdots n_t$, we show surjectivity. But by definition, the map above sends E_1 to $(1/n_1, 0 \cdots, 0)$ in $(\mathbb{Q}/\mathbb{Z})^t$, hence to one of the obvious generators of $\ker N$. \square

Remark 5.4. According to [7], (5.1), the discriminant group $D(\Gamma)$ can be generated by any collection of $t - 1$ of the classes e_1, \dots, e_t of the leaves. However, $D(\Gamma^*)$ might require more than $t - 1$ generators, and even the t leaf classes might not generate.

Example 5.5. Consider the decorated graph Γ^* below of a D_4 singularity:



Index the three leaves clockwise by e_1, e_2, e_3 , and the central curve by f . Then the discriminant group $D(\Gamma)$ is the direct sum of two cyclic groups of order 2, generated by e_1 and e_2 ; note $e_3 = e_1 + e_2$, and $f = 0$. However, $D(\Gamma^*)$ is the direct sum of two cyclic groups of order 4, generated by e_1 and e_2 , and an additional cyclic group of order 2, generated by f (which is not in the span of e_1, e_2, e_3). So the kernel of the natural projection is $\langle 2e_1 \rangle \oplus \langle 2e_2 \rangle \oplus \langle f \rangle$.

On the other hand, we have the following

Proposition 5.6. *The orbifold discriminant group $D(\Gamma^*)$ requires at most t generators.*

Proof. Let $s = m - t$ denote the number of interior vertices of Γ . We claim that the second set of s relations in (5.2) imply that s of the e_i can be written as combinations of the t remaining e_j , which then suffice as generators of $D(\Gamma^*)$.

We use the notation e_i to denote the corresponding vertex of Γ . An interior vertex e_k of valence r gives rise to a relation $-(E_k \cdot E_k)e_k + e_{k,1} + \cdots + e_{k,r} = 0$, so that any $e_{k,j}$ can be written as a combination of the r other e 's. In other words, if e_j is any neighbor of e_k , then e_j can be written as a combination of e_k and e_k 's remaining neighbors.

Choose any interior vertex f , which we label the ‘‘center.’’ We will connect every adjacent pair of vertices with a red or green arrow pointing away from f . First, insert a red arrow from f to one of its neighbors f' . From f to any of its other neighbors insert a green arrow. Next, if the end of an arrow is an end of Γ , do nothing more at that vertex. Otherwise, at any vertex which is the end of an arrow, choose a red arrow to one of its (other) neighbors; if there are any further neighbors, choose a green arrow to each of them. Continue until one reaches all the end vertices of Γ . In this way, for every interior vertex, there is one red arrow emerging. Call the s vertices which are the end of red arrows ‘‘red vertices’’, while the remainder (there are t of them) are ‘‘green vertices.’’

We claim that each red vertex can each be written as a combination of the green ones. Every vertex has a distance from the center, so we prove the result by induction on the distance. We have already said that the center v is green, as are all but one of its neighbors. In particular, its red neighbor can be written as a combination of v and v 's green neighbors. Next, let e be any

red vertex. Then it is the end of a red arrow emanating from a vertex e' which is closer to the center. Then e is a combination of e' and e' 's other neighbors. The other neighbors of e' are e'' , which is closer to the center, and possibly extra ones (if the valence of e' is at least 3). But those extra neighbors are green, since only one red arrow can emerge from e' , and that is the one to e . Thus, e can be written as a combination of green vertices plus e' and e'' . But e' and e'' are closer to the center, so the inductive step applies (whether or not either is red or green). \square

6. ORBIFOLD SPLICE QUOTIENTS

We briefly outline the basics of splice quotient singularities; see Section 1 of [8] for a few more details, and [7] for the full story.

We retain the same notation as before for a good resolution of a singularity, including the graph Γ and the diagonal representation $D(\Gamma) \hookrightarrow (\mathbb{C}^*)^t$. Γ has leaves (vertices of valence 1) and nodes (vertices of valence ≥ 3). Associated to Γ is a *splice diagram* Δ , obtained by collapsing all vertices of valence two, and assigning a positive integer weight to every node and emanating edge by taking an appropriate determinant. From this data, for each node one can assign a weight to every leaf of Δ ; the *semigroup condition* on Δ then requires that each weight on an emanating edge of that node is in the semigroup generated by the weights of the ‘‘outer’’ leaves. Choosing a coordinate x_i for each of the t leaves, one can then write a total of $t - 2$ *splice diagram equations* in the x_i 's, by writing for each node a sum of monomials (one for each emanating edge) which have the same weight as the product of weights around the node (one uses the coefficients of the semigroup relations). The *congruence condition* allows one to insure that each equation so obtained transforms by a character under the action of the representation of $D(\Gamma)$ on \mathbb{C}^t . One thus produces from Γ isolated complete intersection singularities in \mathbb{C}^t on which $D(\Gamma)$ acts, freely off the origin. There are choices involved, and higher order terms can be added.

Γ is called *quasi-minimal* if any string in the graph contains no -1 vertex, or consists of a unique -1 vertex. (A *string* is a connected subgraph containing no nodes.) The Main Theorem 7.2 of [7] starts with a graph Γ :

Theorem 6.1. [7] *Suppose Γ is quasi-minimal and satisfies the semigroup and congruence conditions. Then:*

- (1) *A set of splice diagram equations $\{f_j(x_i) = 0\}$ for Γ defines an isolated complete intersection singularity $(X', 0) \subset (\mathbb{C}^t, 0)$.*
- (2) *The discriminant group $D(\Gamma)$ acts freely on $X' - \{0\}$.*
- (3) *The quotient $(X, 0) \equiv (X', 0)/D(\Gamma)$ has an isolated normal surface singularity, and a good resolution $(\tilde{X}, E) \rightarrow (X, 0)$ whose associated dual graph is Γ .*
- (4) *$f : (X', 0) \rightarrow (X, 0)$ is the universal abelian cover, in particular unramified off the origins.*
- (5) *Each curve $C'_i = \{x_i = 0\} \cap X'$ is mapped by $(X', 0) \rightarrow (X, 0)$ to an irreducible curve C_i , whose proper transform \tilde{C}_i on \tilde{X} is smooth and intersects the exceptional curve transversally, along an end E_i .*

Such $(X, 0)$ are called *splice quotient singularities*.

In more detail, the curve C'_i is reduced, with $|D(\Gamma)|/h$ irreducible components, where h is the order of e_i in $D(\Gamma)$ (follows from [7], Section 3). x_i^h is $D(\Gamma)$ invariant, hence a function on $(X, 0)$; its zero-set is a Cartier divisor, which is h times an irreducible Weil divisor C_i . x_i^h is an *end-curve function* for $(X, 0)$; on the resolution (\tilde{X}, E) it vanishes only on E and an *end-curve* \tilde{C}_i , which intersects E_i transversally at one point. Thus the above construction gives not only a splice-quotient singularity $(X, 0)$, but also a collection of end-curve functions and their corresponding

irreducible curves C_i . Note finally that $f^*(C_i)$ is the reduced sum of the irreducible components of C'_i .

Whether a given $(X, 0)$ is a splice quotient singularity is given by the principal result of [8], the *End-Curve Theorem*:

Theorem 6.2. [8] *Let $(X, 0)$ be a normal surface singularity with $\mathbb{Q}HS$ link, $(\tilde{X}, E) \rightarrow (X, 0)$ a good resolution. Suppose for every end E_i of the exceptional curve E there is a function on $(X, 0)$ whose zero set on \tilde{X} is an end-curve for E_i . Then $(X, 0)$ is a splice quotient singularity.*

An immediate Corollary is the result of T. Okuma:

Theorem 6.3. [10] *Let $(\tilde{X}, E) \rightarrow (X, 0)$ be a quasi-minimal resolution of a rational surface singularity. Then the graph Γ satisfies the semigroup and congruence conditions. Moreover, every end-curve \tilde{C}_i is cut out by an end-curve function on X .*

Proof. For any end-curve \tilde{C}_i on \tilde{X} , the image C_i on X is a \mathbb{Q} -Cartier divisor (since X is rational), so some multiple of it is the zero-set of a function on X . □

To analyze a pair (X, C) , one needs a well-adapted log resolution.

Proposition 6.4. *A pair $(X, C = \sum n_i C_i)$ has a minimal orbifold resolution, a smallest log resolution $\pi : (\tilde{X}_C, E) \rightarrow (X, 0)$ satisfying*

- (1) *each C_i has proper transform \tilde{C}_i which is smooth, intersecting E transversally*
- (2) *\tilde{C}_i intersects a leaf E_i of E*
- (3) *each leaf E_j of E intersects at most one \tilde{C}_i*
- (4) *\tilde{X}_C is quasi-minimal.*

\tilde{X}_C is unique except when a log resolution consists of a single curve plus two \tilde{C}_i intersecting it transversally at two points.

Proof. Starting with the minimal good resolution of X , resolve in a minimal way the singularities of the reduced inverse image of each C_i until one has strong normal crossings (the minimal log-resolution). If the proper transform of some C_i does not intersect a leaf, blow-up that intersection point. If three or more proper transforms intersect the same leaf, blow-up each of the intersection points. If two intersect the same leaf, which has another connection with the graph, again blow-up those points. The only missing case is when two transforms intersect a graph consisting of a single curve; in that case, one could blow-up either of the intersection points. Otherwise, one has uniqueness of the resolution.

We claim the corresponding resolution is quasi-minimal. If a -1 curve intersects some \tilde{C}_i , then it is an end-curve; if its neighboring curve had valence 2, the -1 curve could be blown down, contradicting the minimality of the blow-up process. If a -1 curve intersects no \tilde{C}_i and it has a neighbor curve with valence 2, again one could contract the -1 curve, violating the minimality process. □

We exclude the simple cases for which one has non-uniqueness of the resolution.

Thus, a pair gives rise to a quasi-minimal graph Γ_C , as well as a decorated version Γ_C^* which has arrows and weights at special edges. From this data, we can assign weight one to non-special edges, and encode all the weights in a power map $N : \mathbb{C}^t \rightarrow \mathbb{C}^t$ given on functions by $x_i \mapsto z_i^{n_i}$. By (5.3), one has $D(\Gamma^*) = N^{-1}(D(\Gamma))$.

We give conditions on graphs Γ and Γ^* that allow one to give equations for a pair (X, C) and its UALC. This is Theorem 2 of the introduction, which we now prove.

Let Γ be a quasi-minimal graph with t ends satisfying the semigroup and congruence conditions, and Γ^* a decorated version with arrows and weights at special ends. Assign a coordinate x_i to each end of Γ . Choose a complete intersection singularity $(X', 0) \subset (\mathbb{C}^t, 0)$ defined by $t - 2$ splice equations $\{f_k(x_1, x_2, \dots, x_t) = 0\}$ on which $D(\Gamma)$ acts, freely off 0. Then $(X', 0) \rightarrow (X', 0)/D(\Gamma) \equiv (X, 0)$ is the UAC, and there is a resolution $(\tilde{X}, E) \rightarrow (X, 0)$ with graph Γ . If E_i is a special end of E , the curve $C'_i = \{x_i = 0\} \cap X'$ is reduced and maps to a \mathbb{Q} -Gorenstein curve C_i on $(X, 0)$. Let $C' = \Sigma C'_i$. Consider as above a power map $N : \mathbb{C}^t \rightarrow \mathbb{C}^t$.

Theorem 6.5. *Consider $\Gamma, \Gamma^*, (X', 0), f_k(x_1, \dots, x_t), (X, 0), C_i, C', N : \mathbb{C}^t \rightarrow \mathbb{C}^t$, as above.*

- (1) *Define $(X'', 0) = N^{-1}(X', 0) = \{f_k(z_1^{n_1}, \dots, z_t^{n_t}) = 0, 1 \leq k \leq t - 2\} \subset \mathbb{C}^t$. Then $(X'', 0)$ is an isolated complete intersection singularity on which $D(\Gamma^*)$ acts.*
- (2) *Let $C'' \subset X''$ be the reduced Cartier divisor which is the sum of $\{z_i = 0\} \cap X''$ for special i . Then $D(\Gamma^*)$ acts on C'' and acts freely on $X'' - C''$.*
- (3) *The quotient $(X'', C'') \rightarrow (X'', C'')/D(\Gamma^*) = (X, C)$ is the universal abelian log cover, where $C = \Sigma n_i C_i$ is the sum (with the weights) corresponding to the special ends of Γ^* .*

Proof. We show first that $(X'', 0)$ is non-singular away from the origin. N is a covering map off the intersection with the coordinate hyperplanes $z_i = 0$. N can be factored as the composition of maps which raise powers one coordinate at a time. To study what happens over the reduced curve $\{x_i = 0\} \cap X'$, factor the map N by first raising the i^{th} coordinate to the n_i^{th} power, and then raising all the other coordinates to the appropriate power. The inverse image of $\{x_i = 0\} \cap X'$ under the first map is smooth (away from 0), and the second map is unramified over the new curve. Further, $\{z_i = 0\} \cap X''$ is reduced.

Note $D(\Gamma^*)$ is a subgroup of $(\mathbb{C}^*)^t$, which itself preserves all $\{z_j = 0\}$, so it acts on X'' as well as C'' . By construction, $N : (X'', 0) \rightarrow (X', 0)$ is a covering map off C'' . Since $D(\Gamma)$ acts freely on $X' - \{0\}$, then $N^{-1}(D(\Gamma)) = D(\Gamma^*)$ acts freely on $X'' - C''$. The full (and free) quotient of $X'' - C''$ by $D(\Gamma^*)$ is thus $X - \Sigma C_i$. The construction above matches that of the UALC as described in Section 3. \square

Definition 6.6. A pair which arises as in the Theorem is called an *orbifold splice quotient*.

Remark 6.7. Note that the condition for $(X, \Sigma n_i C_i)$ to be a splice quotient is independent of the weights n_i , and depends only on the C_i (which must be \mathbb{Q} -Cartier).

There is an analogue of the End-Curve Theorem for orbifold splice quotients.

Theorem 6.8. *Let $(X, C = \Sigma n_i C_i)$ be a pair, with minimal orbifold resolution and data (\tilde{X}, E) , Γ, Γ^* . Then the following are equivalent:*

- (1) *(X, C) is an orbifold splice quotient*
- (2) *Each C_i is \mathbb{Q} -Cartier, and for every non-special end E_j of E there is an end-curve function.*

Proof. One implication follows from the above discussion; we prove the converse.

Since C_i is \mathbb{Q} -Cartier, it is the zero-locus of a function g_i on X . Therefore, on \tilde{X} , g_i vanishes only on E and \tilde{C}_i , hence is an end-curve function for that special leaf. But by assumption, for each non-special leaf there is an end-curve function. Thus, there are end-curve functions for every end of Γ . By the End-Curve Theorem, Γ satisfies the semigroup and congruence conditions and $(X, 0)$ itself is a splice quotient. It follows from the construction above that (X, C) is an orbifold splice quotient. \square

Corollary 6.9. *If $(X, 0)$ has a rational singularity, then any pair (X, C) is an orbifold splice quotient.*

Proof. Since $(X, 0)$ is rational, every curve C_i is \mathbb{Q} -Cartier. For each non-special end E_j on the minimal orbifold resolution \tilde{X} , choose any end-curve \tilde{C}_j . Its image C_j on X is the zero-set of a function h_j , which becomes an end-curve function of E_j . The result now follows from the Theorem. \square

Therefore, if $(X, 0)$ has a rational singularity and $C \subset X$ is a reduced curve, then modulo taking an abelian quotient, one can write down from the orbifold resolution diagram Γ some explicit equations for the singularity and the components of the curve C . This includes the case $X = \mathbb{C}^2$, where it has been known for a long time (e.g. [1]) how to write down the equation of a singular curve from its resolution diagram. One should compare as well with Example 3.5.

Corollary 6.10. *Consider the pair $(X, \Sigma_n C_i)$. Suppose $(X, 0)$ is a splice quotient, each C_i is \mathbb{Q} -Cartier, and all the non-special ends of E on the minimal orbifold resolution are ends from the minimal good resolution. Then (X, C) is an orbifold splice quotient.*


Proof. Since $(X, 0)$ is a splice quotient, there is an end-curve function for every end of the graph of the minimal good resolution. Such a function works just as well if that end is also an end on the minimal orbifold resolution. Now apply the Theorem. \square

If $(X, 0)$ is a splice-quotient singularity which is not rational, there will likely exist quasi-minimal resolutions for which the semigroup conditions are not satisfied.

Example 6.11. The minimally elliptic hypersurface singularity $(X, 0) = \{x^2 + y^3 + z^7 = 0\}$ is a splice quotient, with graph and splice diagram for the minimal good resolution



Blowing up further gives a quasi-minimal resolution graph and splice diagram:




The splice diagram does not satisfy the semigroup conditions.

One can understand the previous example as follows: A singularity with this graph (there are two of them) is minimally elliptic and has unimodular resolution graph. Thus the divisor class group is isomorphic to \mathbb{C} , so there is no torsion, and \mathbb{Q} -Cartier is the same as Cartier. The only end-curve functions for the MGR on the -7 curve are powers of z ; so on the further blow-up, one cannot have end-curve functions for both the new -1 end-curves.

7. AN EXAMPLE

Let $(X, 0)$ be the D_4 singularity $\{w^2 = xy(x + y)\}$. Consider the curve $C = pC_1 + qC_2$, where C_1 is the Weil divisor given by the prime ideal (y, w) , and C_2 is a curve, equations to be determined, which on the minimal resolution is smooth and intersects tangentially the central curve, to order 2, away from the other intersection points. To go from the familiar minimal resolution of X to an orbifold resolution, one needs to blow up three times so that \tilde{C}_2 intersects a unique leaf. The graph Γ_C^* is



The representation on \mathbb{C}^5 of $D(\Gamma)$ is the four-group, acting only on the x_1, x_2, x_3 coordinates, by multiplying two of the variables by -1 . There will be 2 splice diagram equations coming from the valence 4 node (locating the cross-ratio of the fourth point), and one from the other. One set of equations (there are many) on which the group acts equivariantly is

$$\begin{aligned} x_4 - x_2^2 - x_3^2 &= 0 \\ x_1^2 + x_2^2 - x_3^2 &= 0 \\ x_5 - x_4^2 - x_1x_2x_3(x_1^2 + x_2^2 + x_3^2) &= 0. \end{aligned}$$

Invariants for the group action are $x = x_1^2, y = x_2^2, z = x_3^2, w = x_1x_2x_3$, and x_4 and x_5 , and these are related by

$$x_4 = y + z; \quad x + y - z = 0; \quad x_5 = x_4^2 + w(x + y + z).$$

Using $xyz = w^2$, the quotient is the familiar equation $w^2 = xy(x + y)$. The curve C_2 turns out to be a principal (i.e., Cartier) divisor defined by $x_5 = (x + 2y)^2 + 2w(x + y)$. $x_2 = 0$ defines C'_1 , so $x_2^2 = y$ is an end-curve function.

To pass to the UALC, in all equations one replaces x_2 by z_2^p , x_5 by z_5^q , and x_1, x_3, x_4 by z_1, z_3, z_4 respectively. The orbifold homology group in $(\mathbb{C}^*)^5$ is generated by

$$(1/2)[1, 0, 1, 0, 0], \quad (1/2p)[p, 1, 0, 0, 0], \quad \text{and} \quad (1/q)[0, 0, 0, 0, 1],$$

hence is the product of three cyclic groups of orders 2, $2p$, and q , respectively. Here we have used the familiar notation $(1/n)[q_1, \dots, q_k]$ to denote the diagonal group generated by $(\zeta^{q_1}, \dots, \zeta^{q_k})$, where ζ is a primitive n^{th} root of 1.

REFERENCES

- [1] D. Eisenbud and W. D. Neumann, *Three-dimensional link theory and invariants of plane curve singularities*. Ann. Math. Stud. **110**, Princeton. Princeton Univ. Press (1985). DOI: [10.1515/9781400881925](https://doi.org/10.1515/9781400881925)
- [2] J. Kollár, *Singularities of the minimal model program*. Cambridge Tracts in Mathematics **200**. Cambridge: Cambridge Univ. Press (2013).
- [3] D. Mumford, The topology of normal singularities of an algebraic surface and a criterion for simplicity, Publ. Math. IHES **9** (1961), 5-22.
- [4] A. Némethi and T. Okuma, On the Casson invariant conjecture of Neumann-Wahl, J. Algebraic Geometry **18** (2009), no. 1, 135-149. DOI: [10.1090/s1056-3911-08-00493-1](https://doi.org/10.1090/s1056-3911-08-00493-1)

- [5] A. Némethi and T. Okuma, The Seiberg-Witten invariant conjecture for splice-quotients, *J. London Math. Soc.*(2) **78** (2008), 143-154. DOI: [10.1112/jlms/jdn020](https://doi.org/10.1112/jlms/jdn020)
- [6] W.D. Neumann, Notes on Geometry and 3-Manifolds, *Low Dimensional Topology*, Böröczky, Neumann, Stipsicz, eds., Bolyai Society Mathematical Studies **8** (1999), 191–267.
- [7] W. D. Neumann and J. Wahl, Complete intersection singularities of splice type as universal abelian covers, *Geom. Topol.* **9** (2005), 699-755. DOI: [10.2140/gt.2005.9.699](https://doi.org/10.2140/gt.2005.9.699)
- [8] W. D. Neumann and J. Wahl, The End Curve Theorem for normal complex surface singularities, *J. Eur. Math. Soc.* **12** (2010), 471-503. DOI: [10.4171/jems/206](https://doi.org/10.4171/jems/206)
- [9] T. Okuma, Universal abelian covers of rational surface singularities, *J. London Math. Soc.* (2) **70** (2004), 307-324. DOI: [10.1112/s0024610704005642](https://doi.org/10.1112/s0024610704005642)
- [10] T. Okuma, Universal abelian covers of certain surface singularities, *Math. Ann.* **334** (2006), 753-773. DOI: [10.1007/s00208-005-0693-8](https://doi.org/10.1007/s00208-005-0693-8)
- [11] T. Okuma, Another proof of the end curve theorem for normal surface singularities, *J. Math. Soc. Japan* **62** (2010), no. 2, 1-11. DOI: [10.2969/jmsj/06210001](https://doi.org/10.2969/jmsj/06210001)
- [12] H.M. Pedersen, Splice diagram singularities and the universal abelian cover of graph orbifolds, *Geom. Dedicata* **195** (2018), 283-305. DOI: [10.1007/s10711-017-0290-5](https://doi.org/10.1007/s10711-017-0290-5)
- [13] P. Scott, The geometries of 3-manifolds, *Bull. London Math. Soc.* **15** (1983), 401-487.
- [14] W. Thurston, Geometry and topology of three-manifolds. [2002 notes at MSRI](#).

WALTER D. NEUMANN, DEPARTMENT OF MATHEMATICS, BARNARD COLLEGE, COLUMBIA UNIVERSITY, NEW YORK, NY 10027

Email address: neumann@math.columbia.edu

JONATHAN WAHL, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27599-3250

Email address: jmwahl@email.unc.edu